Extremely metal-poor stars in classical dwarf spheroidal galaxies

Tafelmeyer, M.; Jablonka, P.; Hill, V.; Shetrone, M.; Tolstoy, E.; Irwin, M. J.; Battaglia, G.; Helmi, A.; Starkenburg, E.; Venn, K. A.

Published in:
Astronomy & astrophysics

DOI:
10.1051/0004-6361/201014733e

Publication date:
2011

A&A 527, C1 (2011)
DOI: 10.1051/0004-6361/201014733e

Astronomy & Astrophysics

Extremely metal-poor stars in classical dwarf spheroidal galaxies: Fornax, Sculptor, and Sextans (*Corrigendum*)

M. Tafelmeyer[1], P. Jablonka[1,2], V. Hill[3], M. Shetrone[4], E. Tolstoy[5], M. J. Irwin[6], G. Battaglia[7], A. Helmi[5], E. Starkenburg[5], K. A. Venn[8], T. Abel[9], P. Francois[2], A. Kaufer[10], P. North[1], F. Primas[7], and T. Szeifert[10]

[1] Laboratoire d'Astrophysique, École Polytechnique Fédérale de Lausanne (EPFL), Observatoire, 1290 Sauverny, Switzerland
e-mail: pascale.jablonka@epfl.ch
[2] GEPI, Observatoire de Paris, CNRS UMR 8111, Université Paris Diderot, 92125 Meudon Cedex, France
[3] Department Cassiopée, University of Nice Sophia-Antipolis, Observatoire de Côte d'Azur, CNRS, 06304 Nice Cedex 4, France
[4] McDonald Observatory, University of Texas, Fort Davis, TX 79734, USA
[5] Kapteyn Astronomical Institute, University of Groningen, PO Box 800, 9700AV Groningen, The Netherlands
[6] Institute of Astronomy, University of Cambridge, Madingley Road, Cambridge CB3 0HA, UK
[7] European Southern Observatory, Karl-Schwarzschild-str. 2, 85748 Garching bei München, Germany
[8] Dept. of Physics & Astronomy, University of Victoria, 3800 Finerty Road, Victoria, BC V8P 1A1, Canada
[9] Kavli Institute for Particle-Astrophysics and Cosmology, Stanford University, SLAC National Accelerator Laboratory, Menlo Park 94025, USA
[10] European Southern Observatory, Alonso de Cordova 3107, Santiago, Chile

A&A, 524, A58 (2010), DOI: 10.1051/0004-6361/201014733

Key words. stars: abundances – Galaxy: evolution – Galaxy: stellar content – Galaxies: star formation – errata, addenda

This erratum corrects a few typographical errors in the coordinates of some of our metal-poor stars.

Table 1. Log of observations. The exposure times are given for the blue and red arms of UVES.

ID	α (J2000)	δ (J2000)	$t_{\rm blue}$(s)	$t_{\rm red}$(s)	S/N			[Fe/H]$_{\rm CaT}$
					4000 Å	5300 Å	6300 Å	
Sex24-72	10 15 02.65	−01 29 55.9	7215	7215	11	36	49	−2.70
Sex11-04	10 13 41.77	−02 11 24.1	5644	8049	11	38	50	−2.56
Fnx05-42	02 41 30.96	−33 55 44.9	21035	21035	13	34	45	−2.76
Fnx M dwarf	02 40 20.64	−34 12 42.7						
Scl07-49	01 00 05.02	−34 01 16.6		20730		36	41	−2.77
Scl07-50	01 00 01.14	−33 59 21.4	36060	27045	27	30	37	−2.83

Notes. The signal-to-noise ratios are measured at 4000 Å (with the exception of Scl07-050, for which S/N per pixel was measured at 4500 Å), 5300 Å, and 6300 Å. The metallicity estimates are derived from CaT or from an HET/HRS spectrum in the case of Sex24-72.